

Instruction for use

Intibio A2MG PLUS (4ml or 6ml)

Product composition

4ml or 6ml of equine leukocyte and erythrocyte poor plasma containing physiological levels of blood proteins.

Intended Use

Intibio A2MG Plus is an animal derivative, intended for supplementation in cases of deficiency. A veterinary clinical follow-up is recommended 2 to 3 weeks after application.

Product Appearance

A2MG PLUS is presented in a 4ml or 6ml tube with blue cap closure. The blue cap needs to be removed before aspiration of the product through the pierceable membrane.

How is A2MG Plus Prepared?

A patented A2MG filter system removes pro-inflammatory proteins from the blood. An extra incubation step at 37°C activates several proteins and intracellular energy mechanisms.

Each batch is tested for purity (total protein, albumin), impurities (white blood cells, red blood cells), sterility (aerobic bacteria, anaerobic bacteria, fungi and yeast), conformity (colour, label inspection) and infectious diseases (over 20 different pathogens).

Storage Instructions

A2MG PLUS is delivered as a 4ml or 6ml tube and can be stored at:

- ◆ 4°C for 4 weeks
- ◆ -20°C (normal freezer) OR -80°C (specialist freezer) OR in liquid nitrogen (in the gas phase) for 24 months after production date

A2MG PLUS can be refrozen once in an upright position to avoid liquid in the cap of the tube.

Handling Instructions

When stored frozen, the product needs to be thawed in the palm of your hand or a warm water bath and heated to 37°C (body temperature) before use to ensure proper protein unfolding and removal of crystallization after freezing. Swirl the tube to dissolve the blood protein clusters and crystals.

When stored at 4°C, the product needs to be heated to body temperature before application to ensure proper protein unfolding. Swirl the tube to dissolve the blood protein clusters.

Please make sure the tube cap remains closed at all times and check the content for a clear, yellow 4ml or 6ml solution per tube before and after aspiration. In case where precipitations are noticed, make sure it is properly heated (37°C) and swirl the tube before aspiration.

Instruction for use

Intibio A2MG PLUS (4ml or 6ml)

Administration Instructions

Use sterile gloves and aseptically prepare the application site prior to use.

Disposal Instructions

A2MG PLUS needs to be disposed as medical waste.

Additional Information/warnings

- ◆ When the content is not transparent or yellow or when the tube or cap is damaged, please do not apply the product and contact the operator.
- ◆ Do not mix with other products from animal origin or from different donors.
- ◆ When biological products from different donors are used in the same region a wash-out period of 4 weeks needs to be respected.
- ◆ Do not add more volume or other products in the tube as too much pressure may result in product leakage.
- ◆ Intended for veterinary use only.

